Communications in Mathematics **34** (2026), no. 2, Paper no. 1 DOI: https://doi.org/10.46298/cm.15764 ©2026 Irene Gonzálvez, Alfredo Miranda, Julio D. Rossi and Jorge Ruiz-Cases This is an open access article licensed under the CC BY-SA 4.0

A two-player zero-sum probabilistic game that approxi mates the mean curvature flow

Irene Gonzálvez, Alfredo Miranda, Julio D. Rossi and Jorge Ruiz-Cases

Abstract. In this paper we introduce a new two-player zero-sum game whose value function approximates the level set formulation for the geometric evolution by mean curvature of a hypersurface. In our approach the game is played with symmetric rules for the two players and probability theory is involved (the game is not deterministic).

Contents

	_
1.1 Description of the main goal	2
1.1.1 The movement by mean curvature of a hypersurface and its as	soci-
ated elliptic equation	3
1.1.2 A probabilistic game approximation for the elliptic problem .	4
1.2 Statements of the main results	8
1.3 Related results	8
<i>MSC 2020:</i> 53E10, 35D40, 35K65, 91A05	
Keywords: Mean curvature flow, Game theory, Viscosity solutions	
Contact information:	
I. Gonzálvez:	
Affiliation: Department of Mathematics, Autonomous University of Madrid, Spain.	
Email: irene.gonzalvez@uam.es	
A. Miranda:	
<i>Email:</i> amiranda@dm_uba_ar	
J. D. Rossi:	
Affiliation: Department of Mathematics and Statistics, Torcuato Di Tella Universi	ty,
Argentina.	
Email: julio.rossi@utdt.edu	
J. Ruiz-Cases:	
Affiliation: Department of Mathematics, Autonomous University of Madrid, Spain.	
Email: jorge.ruizc@uam.es	

Irene Gonzálvez, Alfredo Miranda, Julio D. Rossi and Jorge Ruiz-Cases

2	Preliminaries	8
	2.1 The mean curvature equation	. 8
	2.1.1 Viscosity solutions	. 9
	2.2 Probability. The Optional Stopping Theorem.	. 11
	2.3 A technical lemma	. 12
_	The value of the game is the unique solution to the DPP	14
3 4	The limit of the value functions for the game is the solution to the me	an
3 4	The limit of the value functions for the game is the solution to the me curvature equation	an 18
3 4	The limit of the value functions for the game is the solution to the me curvature equation 4.1 Uniform bounds	an 18 . 19
3 4	The limit of the value functions for the game is the solution to the merecurvature equation 4.1 Uniform bounds 4.2 Estimates near the boundary	an 18 . 19 . 20
3 4	The limit of the value functions for the game is the solution to the merecurvature equation 4.1 Uniform bounds 4.2 Estimates near the boundary 4.3 The limit PDE	an 18 . 19 . 20 . 22

1 Introduction

1.1 Description of the main goal

The aim of this paper is to study a two-players zero-sum probabilistic game whose value functions approximates the motion by mean curvature of a hypersurface. We show that value functions of the game converge, as a parameter that controls the size of the steps made in the game goes to zero, to the unique viscosity solution to a partial differential equation whose level sets encode the motion by mean curvature.

Game theory was used to obtain a better understanding of the mean curvature flow in [21] and [35]. In those references, the authors considered a deterministic (no probability theory is involved) two-person zero-sum game with asymmetric rules for the players. Our goal here is to introduce a different game in which the two players play with symmetric rules and some randomness is involved. In order to pass to the limit in the value functions of the game and find mean curvature as the limit equation we use viscosity solutions (we will rely on previous theory developed for viscosity solutions to geometric flows, see [8] and [15]).

Motion of a hypersurface by mean curvature is by now well-understood. Its usual interpretation involves the steepest-descent for the perimeter functional. Here we will use a different perspective (using game theory) to study this geometric movement. Convexity is preserved under motion by mean curvature, so the boundary of a convex body shrinks monotonically. Thus, when the initial hypersurface is the boundary of a convex domain, the mean curvature flow can be described in two equivalent ways: by following the moving boundary as an evolving surface, or by specifying for each point inside the domain the arrival time when the moving hypersurface reaches the point. We follow the second interpretation and in our game-theoretic interpretation, the latter viewpoint is associated with a minimum-exit-time problem.

Now, let us describe the main ingredients that we need to state and prove our results. First we introduce an elliptic nonlinear partial differential equation (PDE) related to the

geometric motion by mean curvature and next we describe a game whose value functions approximate solutions to the PDE.

1.1.1 The movement by mean curvature of a hypersurface and its associated elliptic equation

Our aim is to describe how a hypersurface that is the boundary of a connected and strictly convex domain, $S = \partial \Omega_0 \subset \mathbb{R}^N$, $N \ge 2$, evolves according to the mean curvature flow. We will use a level set approach to describe this geometric evolution. Assume that there is a real valued function, u(x), defined for $x \in \Omega_0$, and consider the t superlevel sets of u(x),

$$\Omega_t = \{ x : u(x) > t \}, \quad t \ge 0.$$

In what follows we denote by $\nabla u(x)$ and by $D^2u(x)$ the gradient and the Hessian of u with respect to the spacial variable, x. Assume that $\partial \Omega_t$ is smooth. Take $x \in \partial \Omega_t$ a regular point (x is a regular point if $\nabla u(x) \neq 0$). We have $\nabla u(x) \perp \partial \Omega_t$ and for a unitary vector $v \perp \nabla u(x)$ (notice that v is tangential to the hypersurface $\partial \Omega_t$) the quantity $-\langle D^2 u(x)v, v \rangle$ gives the curvature of $\partial \Omega_t$ in the direction of v. Therefore, under these conditions, the mean of the principal curvatures of $\partial \Omega_t$ at a regular point is given by

$$\kappa = \sum_{i} \kappa_{i} = \operatorname{div}\left(\frac{\nabla u}{|\nabla u|}\right)(x) = \frac{1}{|\nabla u(x)|} \left(\Delta u(x) - \left\langle D^{2}u(x)\frac{\nabla u}{|\nabla u|}(x), \frac{\nabla u}{|\nabla u|}(x)\right\rangle\right).$$

We consider the geometric evolution of the hypersurface $\partial \Omega_t$ moving its points in the direction of the normal vector (pointing inside the set Ω_t) with speed given by the mean curvature, $V = -\kappa$ on $\partial \Omega_t$. Now, we look at the elliptic equation

$$\mathcal{L}(u(x)) := \Delta u(x) - \left\langle D^2 u(x) \frac{\nabla u}{|\nabla u|}(x), \frac{\nabla u}{|\nabla u|}(x) \right\rangle = -1, \tag{1}$$

and its associated Dirichlet problem,

$$\begin{cases} \mathcal{L}(u(x)) = -1, & x \in \Omega_0, \\ u(x) = 0, & x \in \partial \Omega_0. \end{cases}$$
(2)

For this problem it is known that there exists a unique viscosity solution and in addition a comparison principle holds, see [11, 21] and references therein. The analysis of (2) in [11, 21] uses the framework of viscosity solutions. This is necessary because in its classical form the PDE is not well-defined when $\nabla u = 0$. However, the problem has nice solutions. Indeed, for a convex planar domain, the evolution remains smooth under motion by mean curvature, and it becomes asymptotically circular as it shrinks to a point [20]. Concerning regularity, in [21] it is proved that u is $C^3(\Omega_0)$.

Notice that when u solves (2) then v(x,t) = u(x) - t is a solution to $v_t(x,t) = \mathcal{L}(v(x,t))$ that is usually referred to as the mean curvature evolution equation. As we have mentioned, for each point inside the domain, we look for the arrival time of the hypersurface. Remark that $\Omega_t = \{x : v(x,t) > 0\} = \{x : u(x) > t\}$. Therefore, (2) is the PDE associated to

the level set formulation for the first arrival time for the geometric evolution of a convex hypersurface by mean curvature.

The motion by mean curvature is nowadays a classical subject, we refer to [6, 11, 12, 14, 17, 25, 26] and also to [18, 32, 33, 35]. Concerning game theoretical approximations for this problem, we quote [21], where the authors introduce a two-players zero-sum game whose value functions approximate solutions to (2). This game is played with asymmetric rules among the players (one chooses a direction and the other a sign) and does not involve probability (the next position of the game only depends on the choices made by the players). See also [19, 35] for other variants of this game. As we have mentioned, here we introduce a new game in which the two players play with symmetric rules and some randomness is involved in the choice of the next position of the game.

1.1.2 A probabilistic game approximation for the elliptic problem

Next, let us describe a game whose value function approximates the solution to (2).

The game is a probabilistic two-person zero-sum game. As in [21] we use Paul for the name of the first player and Carol for the second player. Take $\varepsilon > 0$ (a parameter that controls the size of the possible movements in the game), $\Omega_0 \subset \mathbb{R}^N$ a strictly convex and bounded domain. Let $x_0 \in \Omega_0$ be the initial position of the game. The game is played as follows: at the *i*-th round, Paul chooses a set of unitary vectors $A_i \subset S^{N-1}$ with surface measure $\sigma(A_i) \geq \frac{1}{2}\sigma(S^{N-1}) + \delta_{\varepsilon}$ (with $\delta_{\varepsilon} \approx \varepsilon^{1/2}$) and Carol chooses a set $B_i \subset S^{N-1}$ with measure $\sigma(B_i) \geq \frac{1}{2}\sigma(S^{N-1}) + \delta_{\varepsilon}$. Here and in what follows we denote by σ the surface measure on the sphere S^{N-1} . Notice that both players choose a set slightly bigger than half of the sphere and therefore the intersection of both sets, $A_i \cap B_i$, has positive measure. Once these choices are made, the next position of the game is given by

$$x_i = x_{i-1} + v_i \varepsilon,$$

where the vector v_i is randomly chosen (with uniform probability) in the set $A_i \cap B_i$. The game ends when the position exits Ω_0 and Carol pays to Paul an amount proportional to the number of plays, that is, the payoff is given by $\varepsilon^2 K \times (\text{number of plays})$ with K a constant that we will specify latter.



Figure 1: Game starting at x_0 .

We define $u^{\varepsilon}(x)$ as the value of this game starting at x (the value of the game is just the expected final payoff optimized by both players, Paul wants to maximize the expected outcome, while Carol aims to minimize it). The value of the game for Paul is given by the following formula,

$$u_p^{\varepsilon}(x_0) = \inf_{S_c} \sup_{S_p} \mathbb{E}_{S_p,S_c}^{x_0} \Big[\varepsilon^2 K \times (\text{number of plays}) \Big].$$

Here the inf and sup are taken among the possible strategies for Paul and Carol, that we denote by S_p and S_c , respectively. This is the best possible outcome that Paul may obtain, provided that both players play their best. Analogously, the value for Carol is given by

$$u_c^{\varepsilon}(x_0) = \sup_{S_p} \inf_{S_c} \mathbb{E}_{S_p,S_c}^{x_0} \Big[\varepsilon^2 K \times (\text{number of plays}) \Big].$$

Notice that we always have $u_p(x_0) \ge u_c(x_0)$ for every $x_0 \in \Omega$. Finally, the value of the game is defined as

$$u^{\varepsilon}(x_0) := u_p^{\varepsilon}(x_0) = u_c^{\varepsilon}(x_0)$$

provided the values for Paul and Carol coincide.

For our game there is a value that is given by the unique solution to

To simplify the notation, each time we compute a supremum or an infimum among sets we understand that we are considering subsets of S^{N-1} with surface measure bigger or equal than $\frac{1}{2}\sigma(S^{N-1}) + \delta_{\varepsilon}$. We use $f_C f(v) d\sigma(v)$ for the average

$$\int_C f(v)d\sigma(v) := \frac{1}{\sigma(C)} \int_C f(v)d\sigma(v).$$

The equation that appears in (3) is known as the Dynamic Programming Principle (DPP) in the game literature, see [28], and reflects the rules of the game. In fact, the equation that appears in (3) shows the outcome after only one round of the game starting from x. Paul wants to maximize the outcome, and he can choose the set A, while Carol aims to minimize, and chooses the set B. Then the outcome or the value after making one move is given by the supremum among Paul's choices of the minimum among Carol's choices of the mean value at the new position (the new position is random inside $A \cap B$) plus $K\varepsilon^2$ (the game counts one extra move). We observe that Paul plays aiming to stay inside Ω_0 for a large number of plays while Carol tries to exit Ω_0 as quickly as possible.

Now, to look for a partial differential equation related to this game, we argue formally (one of the main goals of this paper is to make rigorous what follows). Let us find the asymptotic behaviour as $\varepsilon \approx 0$ of the DPP, (3), evaluated at a smooth function. Assume that for a $C^{2,1}$ function ϕ we have

$$\phi(x) \approx \sup_{A} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} \phi(x + v\varepsilon) d\sigma(v) \right\} + \varepsilon^{2} K, \tag{4}$$

that is,

$$0 \approx \sup_{A} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} (\phi(x + v\varepsilon) - \phi(x)) d\sigma(v) \right\} + \varepsilon^{2} K.$$
(5)

Using that $\phi \in C^2$ and neglecting higher order terms, from a simple Taylor expansion, we arrive to

$$0 \approx \sup_{A} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} \frac{1}{\varepsilon} \langle \nabla \phi(x), v \rangle + \frac{1}{2} \langle D^{2} \phi(x)v, v \rangle) d\sigma(v) \right\} + K.$$
(6)

Now, assuming that $\nabla \phi(x) \neq 0$, the leading term is the one that involves $1/\varepsilon$. Hence, when computing the supremum Paul wants to use a vector v such that $\langle \nabla \phi(x), v \rangle$ is as large as possible, and he may choose $A = \{v : \langle \frac{\nabla \phi(x)}{|\nabla \phi(x)|}, v \rangle \geq -\theta_{\varepsilon}\}$ with θ_{ε} such that $\sigma(A) = \frac{1}{2}\sigma(S^{N-1}) + \delta_{\varepsilon}$ (remark that we have the constraint $\sigma(A) \geq \frac{1}{2}\sigma(S^{N-1}) + \delta_{\varepsilon}$). Notice that since $\delta_{\varepsilon} \approx \varepsilon^{1/2}$ we also have $\theta_{\varepsilon} \approx \varepsilon^{1/2}$, in fact, $\sigma(A) \approx \frac{1}{2}\sigma(S^{N-1}) + C\theta_{\varepsilon}$ for θ_{ε} small. Analogously, a clever choice for Carol could be $B = \{v : \langle \frac{\nabla \phi(x)}{|\nabla \phi(x)|}, v \rangle \leq \theta_{\varepsilon}\}$. With these choices of the sets A and B we get $A \cap B = \{v : -\theta_{\varepsilon} \leq \langle \frac{\nabla \phi(x)}{|\nabla \phi(x)|}, v \rangle \leq \theta_{\varepsilon}\}$ and, in this case, by symmetry, the leading term vanishes, since we have

$$\int_{-\theta_{\varepsilon} \le \langle \frac{\nabla \phi(x)}{|\nabla \phi(x)|}, v \rangle \le \theta_{\varepsilon}} \langle \nabla \phi(x), v \rangle d\sigma(v) = 0.$$

Then, from (6) with the sets A and B described before, we arrive to

$$0 \approx \frac{1}{\sigma(\{v : -\theta_{\varepsilon} \le \langle \frac{\nabla \phi(x)}{|\nabla \phi(x)|}, v \rangle \le \theta_{\varepsilon}\})} \int_{-\theta_{\varepsilon} \le \langle \frac{\nabla \phi(x)}{|\nabla \phi(x)|}, v \rangle \le \theta_{\varepsilon}} \frac{1}{2} \langle D^2 \phi(x) v, v \rangle) d\sigma(v) + K.$$
(7)

Observe that the sets $A \cap B = \{v : -\theta_{\varepsilon} \leq \langle \frac{\nabla \phi(x)}{|\nabla \phi(x)|}, v \rangle \leq \theta_{\varepsilon} \}$ converge to $\{v : \langle \nabla \phi(x), v \rangle = 0 \}$ as $\varepsilon \to 0$ and hence, passing to the limit as $\varepsilon \to 0$, we obtain

$$0 = \frac{1}{\mu(\langle \nabla \phi(x), v \rangle = 0)} \int_{\langle \nabla \phi(x), v \rangle = 0} \frac{1}{2} \langle D^2 \phi(x) v, v \rangle) d\mu(v) + K.$$
(8)

Notice that here μ stands for the surface measure in dimension N-2 (we integrate on the set $\{v \in S^{N-1} : \langle \nabla \phi(x), v \rangle = 0\} = S^{N-2}$). At this point we highlight that a rigorous proof of this whole limit procedure is one of the main difficulties in this article.

Now, we just observe that we can compute the Laplacian of ϕ at x using an orthonormal base in which one of the vectors is in the direction of the gradient. We have, assuming for

simplicity that $\nabla \phi(x)$ points in the direction of v_N ,

$$\frac{1}{\mu(\langle \nabla \phi(x), v \rangle = 0)} \int_{\langle \nabla \phi(x), v \rangle = 0} \frac{1}{2} \langle D^2 \phi(x) v, v \rangle) d\mu(v)$$

= $\sum_{i,j=1}^{N-1} \frac{1}{\mu(\langle \nabla \phi(x), v \rangle = 0)} \int_{\langle \nabla \phi(x), v \rangle = 0} \frac{1}{2} \langle D^2 \phi(x) v_i, v_j \rangle) d\mu(v)$
= $C \sum_{i=1}^{N-1} \frac{\partial^2 \phi}{\partial v_i^2}(x) = C \Delta \phi(x) \mid_{\langle \nabla \phi(x), v \rangle = 0}$

for a constant C given by

$$C = \frac{1}{2\mu(\{\langle e_N, v \rangle = 0\})} \int_{\langle e_N, v \rangle = 0} (v_1)^2 \, d\mu(v).$$

Then, (8) is just

$$0 = C\Delta\phi(x) \mid_{\langle \nabla\phi(x), v \rangle = 0} + K.$$
(9)

Hence, if we choose K = C, we arrive to the mean curvature equation

$$0 = \Delta\phi(x) \mid_{\langle \nabla\phi(x), v \rangle = 0} + 1 = \Delta\phi(x) - \left\langle D^2\phi(x) \frac{\nabla\phi}{|\nabla\phi|}, \frac{\nabla\phi}{|\nabla\phi|} \right\rangle + 1.$$
(10)

For other choices of K > 0 we find the equation associated with the movement by a scalar multiple of the mean curvature.

Therefore, we conclude that the DPP associated to the game, (3), is related to the level set formulation of the mean curvature flow. Then, we expect that the limit as $\varepsilon \to 0$ for the value function of the game converges to the viscosity solution to (2).

To make this argument rigorous we run into several difficulties, First, to use viscosity arguments to test the equation with smooth test functions we need to handle inequalities. When we face an inequality we can fix one of the sets, A or B, as above, and still continue with an inequality, however, in order to pass to the limit, we need estimates that hold regardless the choice of the other set. This fact forces us to prove a delicate technical lemma from geometric measure theory, see Lemma 2.5. On top of this, in the limit we need to obtain the boundary condition u = 0 on $\partial\Omega_0$. Since we always have that the values of the game u^{ε} are positive inside Ω_0 (the game is played at least one time before exiting the domain) we only need an upper bound for $u^{\varepsilon}(x)$ for points x inside Ω_0 that are close to the boundary. The fact that this bound has some uniformity in ε (this is needed since we want to pass to the limit as $\varepsilon \to 0$) relies on the fact that we assumed that Ω_0 is strictly convex (this assumption also appears in [21]).

Finally, we introduce the sets

$$\Omega_t^{\varepsilon} = \{ x : u^{\varepsilon}(x) > t \},\$$

that is, the set of positions in \mathbb{R}^N where the players expect to play more than $\left\lceil \frac{t}{k\varepsilon^2} \right\rceil$ rounds. As a consequence of the previous locally uniform convergence of u^{ε} to u, we have that for each t > 0, when ε is small the set where the value function of the ε -game is greater than t, $\Omega_t^{\varepsilon} = \{x : u^{\varepsilon}(x) > t\}$ is close to the set where the solution to the mean curvature problem is greater than t, $\Omega_t = \{x : u(x) > t\}$. Irene Gonzálvez, Alfredo Miranda, Julio D. Rossi and Jorge Ruiz-Cases

1.2 Statements of the main results

Let us state rigorously the main results that are included in this paper. First, let us describe our results for the game. In Section 3 we include the following results.

Theorem 1.1. There exists a unique solution to (3). Moreover, a comparison principle between super and subsolutions to (3) holds.

With this result at hand we can show that the game has a value that coincides with the solution to (3).

Theorem 1.2. The game has a value u^{ε} that is characterized as the unique solution to (3).

Next, in Section 4 we deal with the limit as $\varepsilon \to 0$ of the value function of the game, u^{ε} . Our main result in this paper is the following:

Theorem 1.3. Let $\{u^{\varepsilon}\}_{\varepsilon>0}$ be a family of value functions of the game inside Ω_0 , that is, for each $\varepsilon > 0$, u^{ε} is the solution to (3).

Then, u^{ε} converges uniformly in $\overline{\Omega}_0$ to u, the unique solution to (2), as $\varepsilon \to 0$.

As a consequence, we can obtain the behaviour of the positivity sets of u^{ε} as $\varepsilon \to 0$.

Corollary 1.4. Consider

$$\Omega_t^{\varepsilon} := \{x : u^{\varepsilon}(x) > t\} \text{ and } \Omega_t := \{x : u(x) > t\}.$$
(11)

Then, for each t > 0, we have that

$$\Omega_t \subset \liminf_{\varepsilon \to 0} \Omega_t^\varepsilon \subset \limsup_{\varepsilon \to 0} \Omega_t^\varepsilon \subset \overline{\Omega}_t.$$

1.3 Related results

Let us end this introduction with extra references of related results in the literature.

The level set method for the study of evolution equations for hypersurfaces was first rigorously analyzed in [11, 12]. This method was applied to various equations including the mean curvature flow equation (with or without an obstacle), see [1, 11, 12]. For general references on level sets formulation of geometric flows we refer to [6, 16, 17, 18, 25, 26, 33] and the book [14].

On the other hand, the relation between game theory and nonlinear PDEs is quite rich and has attracted considerable attention in recent years. We quote [4, 21, 22, 24, 27, 29, 30, 34, 35, 36, 37, 38] and the books [5, 23]. Concerning games for geometric flows, as we have mentioned, our immediate precedent is [21] where the authors analyze a deterministic game related to the mean curvature flow. For the formulation of this game in the presence of an obstacle we refer to [35].

2 Preliminaries

2.1 The mean curvature equation

First, we collect some information for the mean curvature problem (2).

2.1.1 Viscosity solutions

Following [21], let us state the precise definition of being a viscosity solution to our elliptic problem,

$$\begin{cases} \Delta u(x) - \left\langle D^2 u(x) \frac{\nabla u}{|\nabla u|}(x), \frac{\nabla u}{|\nabla u|}(x) \right\rangle = -1, & x \in \Omega_0, \\ u(x) = 0, & x \in \partial \Omega_0. \end{cases}$$
(12)

Definition 2.1. An uppersemicontinuous function \underline{u} is a subsolution to (12) provided that if $\underline{u} - \phi$ has a local maximum at a point $x_0 \in \Omega_0$ for some $\phi \in C^2$, then

$$\Delta\phi(x_0) - \left\langle D^2\phi(x_0) \frac{\nabla\phi}{|\nabla\phi|}(x_0), \frac{\nabla\phi}{|\nabla\phi|}(x_0) \right\rangle \ge -1$$

when $\nabla \phi(x_0) \neq 0$; and

$$\Delta\phi(x_0) - \left\langle D^2\phi(x_0)\eta, \eta \right\rangle \ge -1,$$

for some vector η with $|\eta| \leq 1$; and $\underline{u} \leq 0$ on $\partial \Omega$.

A lower semicontinuous function \overline{u} is a supersolution to (12) provided that if $\overline{u} - \phi$ has a local minimum at a point $x_0 \in \Omega_0$ for some $\phi \in C^2$, then

$$\Delta\phi(x_0) - \left\langle D^2\phi(x_0) \frac{\nabla\phi}{|\nabla\phi|}(x_0), \frac{\nabla\phi}{|\nabla\phi|}(x_0) \right\rangle \le -1$$

when $\nabla \phi(x_0) \neq 0$; and

$$\Delta\phi(x_0) - \left\langle D^2\phi(x_0)\eta, \eta \right\rangle \le -1,$$

for some vector η with $|\eta| \leq 1$; and $\overline{u} \geq 0$ on $\partial \Omega$.

Finally, a function u is a solution to (13) if it is both a super and a subsolution.

We can rewrite (12) as

$$\begin{cases} \frac{1}{\mu(\{v:\langle \nabla u(x),v\rangle=0\})} \int_{\langle \nabla u(x),v\rangle=0} \langle D^2 u(x)v,v\rangle d\mu(v) = -C, & x \in \Omega_0, \\ u(x) = 0, & x \in \partial\Omega_0. \end{cases}$$
(13)

Here

$$C = \frac{1}{2\mu(\{\langle e_N, v \rangle = 0\})} \int_{\langle e_N, v \rangle = 0} (v_1)^2 d\mu(v).$$

For completeness, we state the precise definition of being a viscosity solution to (13) (we remark that (12) and (13) are equivalent problems).

Definition 2.2. An uppersemicontinuous function \underline{u} is a subsolution to (13) provided that if $\underline{u} - \phi$ has a local maximum at a point $x_0 \in \Omega_0$ for some $\phi \in C^2$, then

$$\frac{1}{\mu(\{v: \langle \nabla u(x), v \rangle = 0\})} \int_{\langle \nabla u(x), v \rangle = 0} \frac{1}{2} \langle D^2 \phi(x_0) v, v \rangle d\mu(v) \ge -C$$

when $\nabla \phi(x_0) \neq 0$; and

$$\Delta\phi(x_0) - \langle D^2\phi(x_0)\eta, \eta\rangle \geq -1,$$

for some vector η with $|\eta| \leq 1$; and $\underline{u} \leq 0$ on $\partial \Omega$.

A lower semicontinuous function \overline{u} is a supersolution to (13) provided that if $\overline{u} - \phi$ has a local minimum at a point $x_0 \in \Omega_0$ for some $\phi \in C^2$, then

$$\frac{1}{\mu(\{v: \langle \nabla u(x), v \rangle = 0\})} \int_{\langle \nabla u(x), v \rangle = 0} \frac{1}{2} \langle D^2 \phi(x_0) v, v \rangle d\mu(v) \le -C$$

when $\nabla \phi(x_0) \neq 0$; and

$$\Delta\phi(x_0) - \langle D^2\phi(x_0)\eta, \eta\rangle \rangle \le -1,$$

for some vector η with $|\eta| \leq 1$; and $\overline{u} \geq 0$ on $\partial \Omega$.

Finally, a function u is a solution to (13) if it is both a super and a subsolution.

Just for completeness we include a proof of the fact that both definitions are equivalent.

Lemma 2.3. According to Definition 2.1, a function u is a viscosity supersolution (subsolution) if and only if it is a viscosity supersolution (subsolution) according to Definition 2.2.

Proof. The proof is based on the fact that for a smooth function ϕ with $\nabla \phi(x) \neq 0$ it holds that (here we are assuming for simplicity that $\nabla \phi(x)$ points in the direction of e_N),

$$\begin{aligned} \frac{1}{\mu(\langle \nabla \phi(x), v \rangle = 0)} \int_{\langle \nabla \phi(x), v \rangle = 0} \frac{1}{2} \langle D^2 \phi(x) v, v \rangle d\mu(v) \\ &= \sum_{i,j=1}^{N-1} \frac{1}{\mu(\langle \nabla \phi(x), v \rangle = 0)} \int_{\langle \nabla \phi(x), v \rangle = 0} \frac{1}{2} \frac{\partial^2 \phi}{\partial v_i \partial v_j}(x) v_i, v_j d\mu(v) \\ &= C \sum_{i=1}^{N-1} \frac{\partial^2 \phi}{\partial v_i^2}(x) = C \Delta \phi(x) \mid_{\langle \nabla \phi(x), v \rangle = 0} = C \left(\Delta \phi(x) - \left\langle D^2 \phi(x) \frac{\nabla \phi}{|\nabla \phi|}(x), \frac{\nabla \phi}{|\nabla \phi|}(x) \right\rangle \right) \end{aligned}$$

with

$$C = \frac{1}{2\mu(\{\langle \nabla u(x), v \rangle = 0\})} \int_{\langle e_N, v \rangle = 0} (v_1)^2 d\mu(v).$$

Now, assume that u is a viscosity supersolution according to Definition 2.1 and that ϕ is a smooth function such that $u - \phi$ has a local minimum at a point $x_0 \in \Omega_0$. Further assume that $\nabla \phi(x_0) \neq 0$ (otherwise there is nothing to prove). Then, we have

$$\Delta\phi(x_0) - \left\langle D^2\phi(x_0) \frac{\nabla\phi}{|\nabla\phi|}(x_0), \frac{\nabla\phi}{|\nabla\phi|}(x_0) \right\rangle \le -1.$$

From our previous computation this means that,

$$\frac{1}{\mu(\{\langle \nabla \phi(x_0), v \rangle = 0\})} \int_{\langle \nabla \phi(x_0), v \rangle = 0} \frac{1}{2} \langle D^2 \phi(x_0) v, v \rangle d\mu(v) \le -C.$$

Conversely, if for a touching test function with non-vanishing gradient we have

$$\frac{1}{\mu(\{\langle \nabla \phi(x_0), v \rangle = 0\})} \int_{\langle \nabla \phi(x_0), v \rangle = 0} \frac{1}{2} \langle D^2 \phi(x_0) v, v \rangle d\mu(v) \le -C,$$

then, using again our previous computation, we get

$$\Delta\phi(x_0) - \left\langle D^2\phi(x_0) \frac{\nabla\phi}{|\nabla\phi|}(x_0), \frac{\nabla\phi}{|\nabla\phi|}(x_0) \right\rangle \le -1.$$

The fact that u is a viscosity subsolution to the problem according to Definition 2.1 if and only if it is a viscosity subsolution according to Definition 2.2 is completely analogous. It can be obtained just reversing the inequalities when appropriate.

Now, we collect some well known results for the elliptic PDE problem (13). First, a comparison principle holds for viscosity sub and supersolutions to (2) and as a consequence we have uniqueness of solutions. Moreover, combining Perron's method with the comparison principle, we obtain also existence of solutions. For the proofs we refer to [11].

Theorem 2.4 (Comparison Principle). If \underline{u} is a viscosity subsolution and \overline{u} a supersolution to problem (2), then

$$\underline{u}(x) \le \overline{u}(x) \qquad \text{ in } \overline{\Omega}_0.$$

(Existence and uniqueness for the PDE) There exists a unique solution u to (2). Moreover, u is a continuous function in $\overline{\Omega}_0$.

Let us point out that when the domain is a large ball $\Omega_0 = B_R(0)$ the solution to

$$\begin{cases} \Delta u(x) - \left\langle D^2 u(x) \frac{\nabla u}{|\nabla u|}(x), \frac{\nabla u}{|\nabla u|}(x) \right\rangle = -L, & x \in \Omega_0, \\ u(x) = 0, & x \in \partial \Omega_0, \end{cases}$$
(14)

is explicit. Computing the Laplacian in (14) in radial coordinates we find that the solution is given by

$$u(x) = \frac{L}{2(N-1)}(R^2 - |x|^2).$$

Hence, taking L > 1 and R large such that $\Omega_0 \subset B_R(0)$ we get a strict supersolution to our problem (12).

We will also need a very useful tool from probability theory.

2.2 Probability. The Optional Stopping Theorem.

We briefly recall (see [39]) that a sequence of random variables $\{M_k\}_{k\geq 1}$ is a supermartingale (submartingales) if

$$\mathbb{E}[M_{k+1}|M_0, M_1, ..., M_k] \le M_k \ (\ge).$$

Then, the Optional Stopping Theorem, that we will call (OSTh) in what follows, says: given τ a stopping time such that one of the following conditions hold,

(a) The stopping time τ is bounded almost surely;

(b) It holds that $\mathbb{E}[\tau] < \infty$ and there exists a constant c > 0 such that

$$\mathbb{E}[M_{k+1} - M_k | M_0, \dots, M_k] \le c_1$$

(c) There exists a constant c > 0 such that $|M_{\min\{\tau,k\}}| \le c$ almost surely for every k. Then,

$$\mathbb{E}[M_{\tau}] \le \mathbb{E}[M_0] \ (\ge)$$

if $\{M_k\}_{k>0}$ is a supermartingale (submartingale).

For the proof of this classical result we refer to [9, 39].

2.3 A technical lemma

Now, we prove a technical lemma that will be the key ingredient to pass to the limit in the DPP and obtain viscosity solutions to the mean curvature equation. In the next statement recall that σ and μ denote the surface measure on the spheres S^{N-1} and S^{N-2} respectively. Here, we use $v = (\tilde{v}, v_N) \in S^{N-1}$ to denote a unitary vector with $\tilde{v} \in \mathbb{R}^{N-1}$ and $v_N := \langle v, e_N \rangle$, the last coordinate of v.

Lemma 2.5. Let $\delta_{\varepsilon} = \varepsilon^{1/2}$. Take two families of sets, $\{B_{\varepsilon}^*\}_{\varepsilon>0} \subset S^{N-1}$ such that

$$B_{\varepsilon}^* = \Big\{ v \in S^{N-1} : \langle e_N, v \rangle \le \theta_{\varepsilon} \Big\},\$$

with θ_{ε} such that $\sigma(B_{\varepsilon}^*) = \frac{1}{2}\sigma(S^{N-1}) + \delta_{\varepsilon}$, and $\{A_{\varepsilon}^*\}_{\varepsilon>0} \subset S^{N-1}$ any family verifying $\sigma(A_{\varepsilon}^*) \geq \frac{1}{2}\sigma(S^{N-1}) + \delta_{\varepsilon}$. Assume that

$$-\varepsilon \leq \frac{1}{\sigma(A_{\varepsilon}^* \cap B_{\varepsilon}^*)} \int_{A_{\varepsilon}^* \cap B_{\varepsilon}^*} \langle e_N, v \rangle d\sigma(v) \leq 0.$$

Then, given a continuous function $f: S^{N-1} \to \mathbb{R}$, it holds that

$$\frac{1}{\sigma(A_{\varepsilon}^{*}\cap B_{\varepsilon}^{*})} \int_{A_{\varepsilon}^{*}\cap B_{\varepsilon}^{*}} f(v)d\sigma(v) \longrightarrow \frac{1}{\mu(S^{N-2})} \int_{S^{N-1}\cap\{\langle e_{N},v\rangle=0\}} f((\widetilde{v},0))d\mu(\widetilde{v})$$
(15)

when $\varepsilon \to 0$.

Proof. First, we notice that since $\delta_{\varepsilon} \approx \varepsilon^{1/2}$ we have $\theta_{\varepsilon} \approx \varepsilon^{1/2}$.

Let us define the set $U_{\varepsilon} = \{v \in S^{N-1} : |\langle e_N, v \rangle| \le \theta_{\varepsilon}\}$ and consider

$$\begin{split} \left| \int_{A_{\varepsilon}^{*} \cap B_{\varepsilon}^{*}} f(v) d\sigma(v) - \int_{S^{N-1} \cap \{\langle e_{N}, v \rangle = 0\}} f((\widetilde{v}, 0)) d\mu(\widetilde{v}) \right| \\ \leq \underbrace{\left| \int_{A_{\varepsilon}^{*} \cap B_{\varepsilon}^{*}} f(v) d\sigma(v) - \int_{U_{\varepsilon}} f(v) d\sigma(v) \right|}_{I} \\ + \underbrace{\left| \int_{U_{\varepsilon}} f(v) d\sigma(v) - \int_{S^{N-1} \cap \{\langle e_{N}, v \rangle = 0\}} f((\widetilde{v}, 0)) d\mu(\widetilde{v}) \right|}_{II}. \end{split}$$

Let us estimate the second term, *II*. It holds that

$$\lim_{\varepsilon \to 0} \frac{\sigma(U_\varepsilon)}{2\theta_\varepsilon} = \mu(S^{N-2}).$$

Using that f is continuous, by Lebesgue differentiation theorem, we have

$$\frac{1}{2\theta_{\varepsilon}} \int_{\{-\theta_{\varepsilon} \le v_N \le \theta_{\varepsilon}\}} f((\widetilde{v}, v_N)) dv_N \longrightarrow f((\widetilde{v}, 0))$$

when $\varepsilon \to 0$. Hence, we obtain

$$\lim_{\varepsilon \to 0} \frac{2\theta_{\varepsilon}}{\sigma(U_{\varepsilon})} \frac{1}{2\theta_{\varepsilon}} \int_{U_{\varepsilon}} f(v) d\sigma(v) = \frac{1}{\mu(S^{N-2})} \int_{S^{N-2}} f((\widetilde{v}, 0)) d\mu((\widetilde{v})).$$

Thus, $II \to 0$ when $\varepsilon \to 0$.

Now, let us consider I, using that $||f||_{\infty} \leq C$, we get

$$\left| \int_{A_{\varepsilon}^{*} \cap B_{\varepsilon}^{*}} f(v) d\sigma(v) - \int_{U_{\varepsilon}} f(v) d\sigma(v) \right| \\\leq \frac{C}{\sigma(U_{\varepsilon})} \Big[\sigma(A_{\varepsilon}^{*} \cap \{ \langle e_{N}, v \rangle < -\theta_{\varepsilon} \}) + \sigma((A_{\varepsilon}^{*})^{c} \cap U_{\varepsilon}) \Big].$$

Here we used that $\sigma(A_{\varepsilon}^* \cap B_{\varepsilon}^*) \geq \sigma(U_{\varepsilon})$. We will prove that

$$\frac{\sigma(A_{\varepsilon}^* \cap \{\langle e_N, v \rangle < -\theta_{\varepsilon}\})}{\sigma(U_{\varepsilon})} \to 0$$

when $\varepsilon \to 0$. Suppose that this is not true, then there exist a positive constant L and a sequence $\varepsilon_j \to 0$ such that

$$\frac{\sigma(A_{\varepsilon_j}^* \cap \{\langle e_N, v \rangle < -\theta_{\varepsilon_j}\})}{\sigma(U_{\varepsilon_j})} \ge L > 0$$

for all $j \in \mathbb{N}$. Then

$$\frac{\sigma(A_{\varepsilon_j}^* \cap \{\langle e_N, v \rangle < -(1+\alpha)\theta_{\varepsilon_j}\})}{\sigma(U_{\varepsilon_j})} \ge \frac{L}{2} > 0,$$

for some $\alpha > 0$ and for all $j \ge j_0$ for some $j_0 \in \mathbb{N}$. From this point, we omit the subindex j for clarity. We can assume that $\sigma(A_{\varepsilon}^* \cap \{-(1+\alpha)\theta_{\varepsilon} < \langle e_N, v \rangle < -\theta_{\varepsilon}\}) = 0$. Let us use

the hypothesis to obtain

$$\begin{split} -\varepsilon &\leq \frac{1}{\sigma(A_{\varepsilon}^{*} \cap B_{\varepsilon}^{*})} \int_{A_{\varepsilon}^{*} \cap B_{\varepsilon}^{*}} \langle e_{N}, v \rangle d\sigma(v) \leq \frac{1}{\sigma(U_{\varepsilon})} \int_{A_{\varepsilon}^{*} \cap B_{\varepsilon}^{*}} \langle e_{N}, v \rangle d\sigma(v) \\ &\leq \frac{1}{\sigma(U_{\varepsilon})} \Big[\int_{A_{\varepsilon}^{*} \cap \{-\theta_{\varepsilon} \leq \langle e_{N}, v \rangle \leq \theta_{\varepsilon}\}} \langle e_{N}, v \rangle d\sigma(v) + \int_{A_{\varepsilon}^{*} \cap \{\langle e_{N}, v \rangle < -(1+\alpha)\theta_{\varepsilon}\}} \langle e_{N}, v \rangle d\sigma(v) \Big] \\ &\leq \delta_{\varepsilon} \frac{\sigma(A_{\varepsilon}^{*} \cap \{\langle e_{N}, v \rangle < -(1+\alpha)\theta_{\varepsilon}\})}{\sigma(U_{\varepsilon})} - (1+\alpha)\theta_{\varepsilon} \frac{\sigma(A_{\varepsilon}^{*} \cap \{\langle e_{N}, v \rangle < -(1+\alpha)\theta_{\varepsilon}\})}{\sigma(U_{\varepsilon})} \\ &\leq -\alpha\theta_{\varepsilon} \frac{\sigma(A_{\varepsilon}^{*} \cap \{\langle e_{N}, v \rangle < -(1+\alpha)\theta_{\varepsilon}\})}{\sigma(U_{\varepsilon})} \leq -\alpha\theta_{\varepsilon} \frac{L}{2} \end{split}$$

which is a contradiction since we have

$$1 \leq -\alpha \frac{L}{2} \frac{\theta_{\varepsilon}}{\varepsilon} \to -\infty, \quad \text{as } \varepsilon \to 0,$$

because $\delta_{\varepsilon} = \varepsilon^{1/2}$ and thus $\theta_{\varepsilon} \approx \varepsilon^{1/2}$. In the previous estimates we used that

$$0 = \int_{U_{\varepsilon}} \langle e_N, v \rangle d\sigma(v) = \int_{A_{\varepsilon}^* \cap \{-\theta_{\varepsilon} \le \langle e_N, v \rangle \le \theta_{\varepsilon}\}} \langle e_N, v \rangle d\sigma(v) + \int_{(A_{\varepsilon}^*)^c \cap \{-\theta_{\varepsilon} \le \langle e_N, v \rangle \le \theta_{\varepsilon}\}} \langle e_N, v \rangle d\sigma(v)$$

implies

$$\frac{1}{\sigma(A_{\varepsilon}^{*}\cap B_{\varepsilon}^{*})} \int_{A_{\varepsilon}^{*}\cap\{-\theta_{\varepsilon}\leq\langle e_{N},v\rangle\leq\theta_{\varepsilon}\}} \langle e_{N},v\rangle d\sigma(v) \\
= \frac{1}{\sigma(A_{\varepsilon}^{*}\cap B_{\varepsilon}^{*})} \int_{(A_{\varepsilon}^{*})^{c}\cap\{-\theta_{\varepsilon}\leq\langle e_{N},v\rangle\leq\theta_{\varepsilon}\}} -\langle e_{N},v\rangle d\sigma(v) \\
\leq \theta_{\varepsilon} \frac{\sigma((A_{\varepsilon}^{*})^{c}\cap\{-\theta_{\varepsilon}\leq\langle e_{N},v\rangle\leq\theta_{\varepsilon}\})}{\sigma(U_{\varepsilon})} \\
= \theta_{\varepsilon} \frac{\sigma(A_{\varepsilon}^{*}\cap\{\langle e_{N},v\rangle<-(1+\alpha)\theta_{\varepsilon}\})}{\sigma(U_{\varepsilon})}.$$

Thus, we also have that $I \to 0$ when $\varepsilon \to 0$. To show that

$$\frac{\sigma((A_{\varepsilon}^*)^c \cap U_{\varepsilon})}{\sigma(U_{\varepsilon})} \to 0$$

is analogous. This ends the proof.

3 The value of the game is the unique solution to the DPP

In this section we show that the value of the game is characterized as the unique solution to the DPP. This fact is crucial when we pass to the limit in the viscosity sense and obtain the limit is a solution to the mean curvature equation.

Lemma 3.1. There exists a solution to

$$\begin{cases} v(x) = \sup_{A \to B} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} v(x + v\varepsilon) d\sigma(v) \right\} + \varepsilon^{2} K, & x \in \Omega_{0}, \\ v(x) = 0, & x \in \mathbb{R}^{N} \setminus \Omega_{0}. \end{cases}$$
(16)

Proof. We start with $w_0 \equiv 0$, and we define inductively

$$w_{n+1}(x) = \sup_{A} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} w_n(x + v\varepsilon) d\sigma(v) \right\} + \varepsilon^2 K, \quad \text{for } x \in \Omega_0$$

with $w_{n+1} \equiv 0$ for $x \in \mathbb{R}^N \setminus \Omega_0$.

First, note that for each x in the interior of Ω_0 , the chosen sets A and B where supremum and infimum are attained are not necessarily contained in Ω_0 . In fact, for points close enough to the boundary, we expect that a most part of the selected B lies outsides Ω_0 . Consequently, w_n is not constant within Ω_0 .

Second, it is not hard to check by an inductive argument that w_n is increasing with n, and thus we can consider

$$v(x) = \lim_{n \to \infty} w_n(x).$$

Notice that

$$u(x) = \frac{L}{2(N-1)}(R^2 - |x|^2).$$

with L > 1 and R large such that $\Omega_0 \subset B_R(0)$ is a strict supersolution to our problem (12). Therefore, by a simple Taylor expansion like the one that we outline in the introduction we have that u(x) is a strict supersolution to (16). This argument provides a uniform upper bound for the sequence w_n .

Now, we observe that, by the monotone convergence theorem we have

$$\lim_{n \to \infty} \sup_{A} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} w_n (x + v\varepsilon) d\sigma(v) \right\} + \varepsilon^2 K$$
$$= \sup_{A} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} v (x + v\varepsilon) d\sigma(v) \right\} + \varepsilon^2 K.$$

Hence, we conclude that

$$v(x) = \sup_{A} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} v(x + v\varepsilon) d\sigma(v) \right\} + \varepsilon^{2} K, \quad \text{for } x \in \Omega_{0}$$

with $v \equiv 0$ for $x \in \mathbb{R}^N \setminus \Omega_0$. Thus, we have the desired existence of a solution to the DPP.

Recall that the value of the game for Paul is given by

$$\overline{u}^{\varepsilon}(x_0) = \inf_{S_c} \sup_{S_p} \mathbb{E}^{x_0}_{S_p,S_c} \Big[\varepsilon^2 K \times (\text{number of plays}) \Big].$$

This is the best value that Carol may guarantee to obtain.

Analogously, the value of the game from Carol's viewpoint is

$$\underline{u}^{\varepsilon}(x_0) = \sup_{S_p} \inf_{S_c} \mathbb{E}^{x_0}_{S_p, S_c} \Big[\varepsilon^2 K \times (\text{number of plays}) \Big].$$

We say that the game has a value if we can reverse inf with sup, that is, when

$$\overline{u}^{\varepsilon}(x_0) = \underline{u}^{\varepsilon}(x_0) := u^{\varepsilon}(x_0).$$

Our next result shows that this game has a value and, moreover, proves that the value function is the solution to the DPP.

Theorem 3.2. The game has a value that is characterized as the unique solution to (16).

Proof. In Lemma 3.1 we proved that the DPP has a solution v(x). In what follows we use this solution to build quasi-optimal strategies for the players and use probabilistic arguments to show that indeed the value of the game coincides with the solution to the DPP, v.

At every position of the game, x_k , Paul (who wants to maximize) chooses A^* in such a way that

$$\sup_{A} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} v(x_{k} + v\varepsilon) d\sigma(v) \right\}$$
$$\leq \inf_{B} \left\{ \frac{1}{\sigma(A^{*} \cap B)} \int_{A^{*} \cap B} v(x_{k} + v\varepsilon) d\sigma(v) \right\} + \frac{\eta}{2^{k+1}}$$

Given this strategy for Paul and any strategy S_c for Carol we consider the sequence of random variables given by

$$M_k = v(x_k) - \frac{\eta}{2^k}$$

Let us see that $(M_k)_{k\geq 0}$ is a submartingale. To this end we need to estimate

 $\mathbb{E}_{S_p^*,S_c}[M_{k+1}\mid_{M_k}].$

Since we are using the strategies S_p^* and S_c , it holds that

$$\mathbb{E}_{S_p^*,S_c}[M_{k+1}\mid_{M_k}] = \mathbb{E}_{S_p^*,S_c}[v(x_{k+1}) - \frac{\eta}{2^{k+1}}\mid_{M_k}]$$
$$= \left\{\frac{1}{\sigma(A^* \cap B)} \int_{A^* \cap B} v(x_k + v\varepsilon) d\sigma(v)\right\} - \frac{\eta}{2^{k+1}}$$

Here A^* corresponds to Paul's choice and B to Carol's. Now, we have

$$\sup_{A} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} v(x_{k} + v\varepsilon) d\sigma(v) \right\}$$

$$\leq \inf_{B} \left\{ \frac{1}{\sigma(A^{*} \cap B)} \int_{A^{*} \cap B} v(x_{k} + v\varepsilon) d\sigma(v) \right\} + \frac{\eta}{2^{k+1}}$$

$$\leq \left\{ \frac{1}{\sigma(A^{*} \cap B)} \int_{A^{*} \cap B} v(x_{k} + v\varepsilon) d\sigma(v) \right\} + \frac{\eta}{2^{k+1}}.$$

Therefore, we arrive to

$$\mathbb{E}_{S_p^*, S_c}[M_{k+1} \mid_{M_k}] \ge \sup_A \inf_B \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} v(x_k + v\varepsilon) d\sigma(v) \right\} - \frac{\eta}{2^k}$$

As v is a solution to the DPP (3) we obtain

$$\mathbb{E}_{S_{p}^{*},S_{c}}[M_{k+1}\mid_{M_{k}}] \ge v(x_{k}) - \frac{\eta}{2^{k}} = M_{k}$$

as we wanted to show.

Therefore, $(M_k)_{k\geq 0}$ is a submartingale. Using the optional stopping theorem (recall that we have that τ is finite a.s. and that we have that M_k is uniformly bounded we conclude that

$$\mathbb{E}_{S_p^*, S_c}[M_\tau] \ge M_0$$

where τ is the first time such that $x_{\tau} \notin \Omega$. Then,

$$\mathbb{E}_{S_p^*, S_c}[\varepsilon^2 K\tau] \ge v(x_0) - \eta.$$

We can compute the infimum in S_c and then the supremum in S_p to obtain

$$\sup_{S_p} \inf_{S_c} \mathbb{E}_{S_p, S_c} [\varepsilon^2 K \tau] \ge v(x_0) - \eta.$$

An analogous computation shows that

$$\inf_{S_c} \sup_{S_p} \mathbb{E}_{S_p, S_c} [\varepsilon^2 K \tau] \le v(x_0) + \eta.$$

To end the proof we just observe that

$$\sup_{S_p} \inf_{S_c} \mathbb{E}_{S_p, S_c} [\varepsilon^2 K \tau] \le \inf_{S_c} \sup_{S_p} \mathbb{E}_{S_p, S_c} [\varepsilon^2 K \tau].$$

Therefore,

$$v(x_0) - \eta \le \sup_{S_p} \inf_{S_c} \mathbb{E}_{S_p, S_c}[\varepsilon^2 K \tau] \le \inf_{S_c} \sup_{S_p} \mathbb{E}_{S_p, S_c}[\varepsilon^2 K \tau] \le v(x_0) + \eta$$

Since $\eta > 0$ is arbitrary, we conclude that the game has a value and that

$$v(x_0) = \sup_{S_p} \inf_{S_c} \mathbb{E}_{S_p, S_c}[\varepsilon^2 K \tau] = \inf_{S_c} \sup_{S_p} \mathbb{E}_{S_p, S_c}[\varepsilon^2 K \tau].$$

That is

$$v(x_0) = \underline{u}^{\varepsilon}(x_0) = \overline{u}^{\varepsilon}(x_0).$$

This ends the proof.

As a consequence of the previous arguments we obtain a comparison principle for sub and super solutions to the DPP. First, let us state what we understand by a super and a subsolution to the DPP.

Definition 3.3. A function $\overline{u}^{\varepsilon}$ is a supersolution to the DPP if it verifies

$$\begin{cases} \overline{u}^{\varepsilon}(x) \ge \sup_{A} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} \overline{u}^{\varepsilon}(x + v\varepsilon) d\sigma(v) \right\} + \varepsilon^{2} K, & x \in \Omega_{0}, \\ \overline{u}^{\varepsilon}(x) \ge 0, & x \in \mathbb{R}^{N} \setminus \Omega_{0}. \end{cases}$$
(17)

A function $\underline{u}^{\varepsilon}$ is a subsolution to the DPP when the reverse inequalities hold, that is, when

$$\begin{cases} \underline{u}^{\varepsilon}(x) \leq \sup_{A} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} \underline{u}^{\varepsilon}(x + v\varepsilon) d\sigma(v) \right\} + \varepsilon^{2} K, & x \in \Omega_{0}, \\ \underline{u}^{\varepsilon}(x) \leq 0, & x \in \mathbb{R}^{N} \setminus \Omega_{0}. \end{cases}$$
(18)

Now, we are ready to prove the comparison argument.

Theorem 3.4. Let $\overline{u}^{\varepsilon}$ be a supersolution to the DPP and \underline{u} a subsolution. Then

$$\overline{u}^{\varepsilon}(x) \geq \underline{u}^{\varepsilon}(x), \qquad \text{for every } x \in \mathbb{R}^N.$$

Proof. We only have to observe that the previous proof shows that $\overline{u}^{\varepsilon}(x)$ is bigger or equal than the value of the game. Indeed, one can follow the same argument as before to show that

$$M_k = \overline{u}^{\varepsilon}(x_k) + \frac{\eta}{2^k}$$

is a supermartingale using that $\overline{u}^{\varepsilon}$ verifies (17).

Analogously, one can also show that \underline{u} is less or equal than the value of the game. Then, we conclude that

$$\overline{u}^{\varepsilon}(x) \ge \underline{u}^{\varepsilon}(x), \qquad \text{for every } x \in \mathbb{R}^N$$

as we wanted to show.

4 The limit of the value functions for the game is the solution to the mean curvature equation

In this section, we will obtain that the unique solution u to (2) is the locally uniform limit of the value functions u^{ε} of the ε -game as ε goes to zero, see Theorem 1.3. Our strategy is to prove that the functions

$$\overline{u}(x) = \limsup_{\substack{\varepsilon \to 0^+ \\ y \to x}} u^{\varepsilon}(y), \quad \underline{u}(x) = \liminf_{\substack{\varepsilon \to 0^+ \\ y \to x}} u^{\varepsilon}(y), \tag{19}$$

are, respectively, a viscosity subsolution and a viscosity supersolution to the problem (2) with $\overline{u}(x) = \underline{u}(x) = 0$ for $x \in \partial \Omega_0$. Therefore, by the Comparison Principle, Theorem 2.4, we get that $\overline{u} \leq \underline{u}$. Since by definition $\underline{u} \leq \overline{u}$, we then obtain that $\overline{u} = \underline{u}$. Hence, the limit of the family u^{ε} exists, and it is the unique solution u to (2). Finally, as a consequence of the convergence $u^{\varepsilon} \to u$, for each t > 0 we will be able to approximate the set where $u(\cdot)$ is bigger than t by the set where $u^{\varepsilon}(\cdot)$ is bigger than t for ε small enough, see Corollary 1.4.

4.1 Uniform bounds

First, we show that the value function of the game, u^{ε} is uniformly bounded.

Lemma 4.1. There exists a constant C, independent of ε , such that

$$\sup_{x \in \Omega_0} |u^{\varepsilon}(x)| \le C.$$
(20)

Proof. We have that $u^{\varepsilon}(x) \geq 0$. Hence, we need to show only an upper bound. To simplify the notation we assume that this point is the origin z = 0. Assume that at any point $x \in \Omega_0$ Carol chooses B_{ε} as the set

$$B_{\varepsilon} = \left\{ v \in S^{N-1} : \langle v, \frac{x}{|x|} \rangle \ge -\theta_{\varepsilon} \right\}$$

with θ_{ε} such that $\sigma(B_{\varepsilon}) = \frac{1}{2}\sigma(S^{N-1}) + \delta_{\varepsilon}$. That is, Carol aims to go as far as possible from z = 0.

Then, for any set A that Paul may choose, we have that the next position of the game provided that we arrive to x_k satisfies

$$\begin{split} \mathbb{E}[|x_{k+1}|^{2}|_{x_{k}}] &= \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A \cap B_{\varepsilon}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \\ &= \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A \cap \{-\theta_{\varepsilon} \leq \langle v, \frac{x_{k}}{|x_{k}|} \rangle \leq \theta_{\varepsilon}\}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \\ &+ \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A \cap \{\theta_{\varepsilon} \leq \langle v, \frac{x_{k}}{|x_{k}|} \rangle \}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \\ &\geq \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A \cap \{-\theta_{\varepsilon} \leq \langle v, \frac{x_{k}}{|x_{k}|} \rangle \leq \theta_{\varepsilon}\}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \\ &+ \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A^{c} \cap \{-\theta_{\varepsilon} \leq \langle v, \frac{x_{k}}{|x_{k}|} \rangle \leq \theta_{\varepsilon}\}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \\ &= \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{\{-\theta_{\varepsilon} \leq \langle v, \frac{x_{k}}{|x_{k}|} \rangle \leq \theta_{\varepsilon}\}} |x_{k} + v\varepsilon|^{2} d\sigma(v) = |x_{k}|^{2} + \varepsilon^{2}. \end{split}$$

Here we used that

$$\sigma(A \cap \{\theta_{\varepsilon} \le \langle v, \frac{x_k}{|x_k|} \rangle\}) \ge \sigma(A^c \cap \{-\theta_{\varepsilon} \le \langle v, \frac{x_k}{|x_k|} \rangle \le \theta_{\varepsilon}\})$$

and

$$x_k + v\varepsilon|^2 \mid_{A \cap \{\theta_\varepsilon \le \langle v, \frac{x_k}{|x_k|} \rangle\}} \ge |x_k + v\varepsilon|^2 \mid_{A^c \cap \{-\theta_\varepsilon \le \langle v, \frac{x_k}{|x_k|} \rangle \le \theta_\varepsilon\}} .$$

Then

$$\frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A \cap \{\theta_{\varepsilon} \le \langle v, \frac{x_{k}}{|x_{k}|} \rangle\}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \ge \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A^{c} \cap \{-\theta_{\varepsilon} \le \langle v, \frac{x_{k}}{|x_{k}|} \rangle \le \theta_{\varepsilon}\}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A^{c} \cap \{-\theta_{\varepsilon} \le \langle v, \frac{x_{k}}{|x_{k}|} \rangle \le \theta_{\varepsilon}\}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A^{c} \cap \{-\theta_{\varepsilon} \le \langle v, \frac{x_{k}}{|x_{k}|} \rangle \le \theta_{\varepsilon}\}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A^{c} \cap \{-\theta_{\varepsilon} \le \langle v, \frac{x_{k}}{|x_{k}|} \rangle \le \theta_{\varepsilon}\}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A^{c} \cap \{-\theta_{\varepsilon} \le \langle v, \frac{x_{k}}{|x_{k}|} \rangle \le \theta_{\varepsilon}\}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A^{c} \cap \{-\theta_{\varepsilon} \le \langle v, \frac{x_{k}}{|x_{k}|} \rangle \le \theta_{\varepsilon}\}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A^{c} \cap \{-\theta_{\varepsilon} \le \langle v, \frac{x_{k}}{|x_{k}|} \rangle \le \theta_{\varepsilon}\}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A^{c} \cap \{-\theta_{\varepsilon} \le \langle v, \frac{x_{k}}{|x_{k}|} \rangle \le \theta_{\varepsilon}\}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A^{c} \cap \{-\theta_{\varepsilon} \le \langle v, \frac{x_{k}}{|x_{k}|} \rangle \le \theta_{\varepsilon}\}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A^{c} \cap \{-\theta_{\varepsilon} \le \langle v, \frac{x_{k}}{|x_{k}|} \rangle \le \theta_{\varepsilon}\}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A^{c} \cap \{-\theta_{\varepsilon} \le \langle v, \frac{x_{k}}{|x_{k}|} \rangle \le \theta_{\varepsilon}\}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A^{c} \cap \{-\theta_{\varepsilon} \le \langle v, \frac{x_{k}}{|x_{k}|} \rangle \le \theta_{\varepsilon}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A^{c} \cap \{-\theta_{\varepsilon} \le \langle v, \frac{x_{k}}{|x_{k}|} \rangle \le \theta_{\varepsilon}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A^{c} \cap \{-\theta_{\varepsilon} \le \langle v, \frac{x_{k}}{|x_{k}|} \rangle \le \theta_{\varepsilon}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} \int_{A^{c} \cap \{-\theta_{\varepsilon} \le \langle v, \frac{x_{k}}{|x_{k}|} \rangle \le \theta_{\varepsilon}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \theta_{\varepsilon}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \theta_{\varepsilon}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \theta_{\varepsilon} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \frac{1}{\sigma(A \cap B_{\varepsilon})} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \theta_{\varepsilon}} |x_{k} + v\varepsilon|^{2} d\sigma(v) \le \theta_{\varepsilon} |x_{k} + v$$

Now, if we introduce the sequence of random variables

$$M_k = |x_k|^2 - \varepsilon^2 k.$$

It holds that

$$\mathbb{E}[M_{k+1}\mid_{M_k}] \ge |x_k|^2 + \varepsilon^2 - \varepsilon^2(k+1) = M_k$$

Then, we have that M_k is a submartingale and using the optional stopping theorem (OSTh), we obtain

$$\mathbb{E}[M_{\tau}] \ge M_0 = |x_0|^2$$

Hence we conclude that

$$\mathbb{E}[\varepsilon^2 \tau] \le C(\Omega)$$

This says that the expected value for the number of plays when Carol uses the previous strategy is bounded by C/ε^2 . Hence, the value of the game verifies

$$u^{\varepsilon}(x_0) \leq \sup_{S_p} \mathbb{E}_{S_p, S_c^*}[\varepsilon^2 \tau] \leq C$$

as we wanted to show.

As an alternative argument, we can use that

$$u(x) = \frac{L}{2(N-1)}(R^2 - |x|^2).$$

with L > 1 and R large such that $\Omega_0 \subset B_R(0)$ is a strict supersolution to our problem (12). Then, by a simple Taylor expansion like the one that we outline in the introduction we have that u(x) is a strict supersolution to (16) for ε small enough. Then, the comparison principle for the DPP, Theorem 3.4, provides a uniform upper bound for the solution to the DPP u^{ε} for ε small enough.

4.2 Estimates near the boundary

Next, we prove that the half-relaxed limits of the family u^{ε} , \overline{u} and \underline{u} , given by (19), attain the boundary condition.

Proposition 4.2. Let \overline{u} and \underline{u} be defined as in (19). Then

$$\overline{u}(x) = \underline{u}(x) = 0,$$
 for every $x \in \partial \Omega_0.$

Proof. We will assume, to simplify the notation, that we start the game at a point x_0 of the form $x_0 = (0, .., 0, x_N^0) \in \Omega_0$ with $x_N^0 > 0$ that is close to $y_0 = 0 \in \partial \Omega_0$.

Now, let us show the two inequalities needed for the proof. The lower bound is straightforward since we have that

$$u_{\varepsilon}(x_0) \ge 0,\tag{21}$$

and then we conclude that

$$\overline{u}(x) \ge \underline{u}(x) \ge 0$$

in the whole $\overline{\Omega}_0$.

To find a reverse estimate is more delicate. To this end we need to choose a strategy for Carol (the player who wants to minimize the expected payoff), associated to the following claim.

CLAIM: Given $\eta > 0$, there exist $r_0 > 0$ and $z \in \mathbb{R}^N$ such that if $|x_0| < r_0$, and $x_{\tau} \in \partial \Omega_0 \cap (B_{|x_0-z|}(z))^c$ for some $\tau \in \mathbb{N}$, we get

$$\left| |x_{\tau} - z|^2 - |x_0 - z|^2 \right| < \eta.$$

Suppose that the claim is true (we will prove it later). At each turn, if the position of the token is in x_k we let Paul choose freely while Carol chooses B^* as the set

$$B^* = \left\{ v \in S^{n-1} : \langle v, \frac{x_k - z}{|x_k - z|} \rangle \ge -\theta_{\varepsilon} \right\},\$$

with θ_{ε} such that $\sigma(B_{\varepsilon}^*) = \frac{1}{2}\sigma(S^{N-1}) + \delta_{\varepsilon}$.

Since Carol aims to minimize the expected payoff, we get a bound from above for the value of the game, it holds that

$$u_{\varepsilon}(x_0) \le \sup_{S_p} \mathbb{E}^{x_0}_{S_c^*, S_p}[\varepsilon^2 \tau].$$
(22)

Then, for any sets A (any strategy for Paul's choices), let us consider the following sequence of random variables

$$M_k = |x_k - z|^2.$$

We have

$$\mathbb{E}[M_{k+1}|_{x_k}] = \mathbb{E}[|x_{k+1} - z|^2|_{x_k}] = \left\{ \frac{1}{\sigma_{n-1}(A \cap B^*)} \int_{A \cap B^*} |x_k - z + \varepsilon v|^2 d\sigma_{n-1}(v) \right\}$$

$$\geq |x_k - z|^2 + \varepsilon^2 = M_k + \varepsilon^2 \geq M_k.$$

Thus, using the optional stopping theorem we get

$$\mathbb{E}[M_{\tau}] \ge M_0,$$

that is

$$\mathbb{E}[|x_{\tau} - z|^2] \ge |x_0 - z|^2.$$

This implies that $x_{\tau} \in \partial \Omega_0 \cap (B_{|x_0-z|}(z))^c$. On the other hand, if we consider

$$N_k = M_k - k\varepsilon^2$$

we obtain

$$\mathbb{E}[N_{k+1}\mid_{x_k}] = \mathbb{E}[M_{k+1} - (k+1)\varepsilon^2\mid_{x_k}] \ge M_k + \varepsilon^2 - (k+1)\varepsilon^2 = N_k$$

that is, N_k is a submartingale. Then, if we use the optional stopping theorem we get

 $\mathbb{E}[N_{\tau}] \ge N_0.$

That is

$$\mathbb{E}[|x_{\tau} - z|^2 - \tau \varepsilon^2] \ge |x_0 - z|^2.$$

Finally, we obtain

$$\mathbb{E}[\tau\varepsilon^2] \le \mathbb{E}[|x_{\tau} - z|^2] - |x_0 - z|^2$$

which is small, using the claim. This implies that

$$\underline{u}(0) = \liminf u_{\varepsilon}(x) = 0 \text{ and } \overline{u}(0) = \limsup u_{\varepsilon}(x) = 0.$$

Let us prove the claim.

Proof of the claim. We can assume that $\overline{\Omega}_0 \subset U$ where

$$U = \left\{ x \in \mathbb{R}^N : x_N \ge c \sum_{j=1}^{N-1} x_j^2 \right\},\$$

with c > 0. Here we use the fact that we assumed that Ω_0 is strictly convex. We can also suppose that $\partial \Omega_0 \cap \partial U = \{0\}$. We will work with the set U. If we consider $z = (0, \ldots, \Lambda)$ with $\Lambda > 0$ large enough. One can check that the function

 $\mu: U \cap \{x: \langle x - x_0, e_N \rangle \le 0\},\$

defined by $\mu(x) = |x - z|^2 - |x_0 - z|^2$ verify $|\mu| < \eta$ if $|x_0|$ is small enough. We can finish the proof of the claim by choosing z such that the set

$$U \cap (B_{|x_0-z|}(z))^c \sim U \cap \{x : \langle x - x_0, e_N \rangle \le 0\}.$$

This ends the proof.

4.3 The limit PDE

Now, let us show that the half-relaxed limits are sub and supersolutions to our elliptic problem.

Theorem 4.3. The functions \overline{u} and \underline{u} defined in (19) are respectively viscosity subsolution and supersolutions to the elliptic problem (2).

Proof. First, remark that \overline{u} is upper semicontinuous and \underline{u} is lower semicontinuous. Let us write the Dynamic Programming Principle of our parabolic game as

 $0 = \sup_{A} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} \left(u^{\varepsilon} (x + v\varepsilon) - u^{\varepsilon}(x) \right) d\sigma(v) \right\} + K\varepsilon^{2}.$

The use of this equation is the key to pass to the limit in the viscosity sense.

Let us show that \overline{u} is a viscosity subsolution to the problem (2). To this end, let $\phi \in C^2$ be a function such as $\phi - \overline{u}$ achieves a strict minimum at x_0 in $\overline{B}_{R_0}(x_0)$ for some $R_0 > 0$ such that $\overline{B}_{R_0}(x_0)$ is in the interior of Ω_0 .

Assume first that $\nabla \phi(x_0) \neq 0$. We want to prove that

$$\Delta\phi(x_0)\mid_{\langle\nabla\phi(x),v\rangle=0}+1\geq 0$$

For every $\varepsilon > 0$, we can choose $x_{\varepsilon} \in \overline{B}_{R_0}(x_0)$ such that

$$\phi(x_{\varepsilon}) - u^{\varepsilon}(x_{\varepsilon}) - \varepsilon^{3} \leq \inf_{\overline{B}_{R_{0}}(x_{0})} \{\phi(x) - u^{\varepsilon}(x)\} \leq \phi(x) - u^{\varepsilon}(x),$$

for every $x \in \overline{B}_{R_0}(x_0)$. As $\overline{B}_{R_0}(x_0)$ is a compact set, x_{ε} converges, up to a subsequence, to some \overline{x} as $\varepsilon \to 0^+$. We have that $\overline{x} = x_0$, since x_0 is the only maximum of $\overline{u} - \phi$ in $B_{R_0}(x_0)$. Rearranging the previous expression, we get

$$u^{\varepsilon}(y) - u^{\varepsilon}(x_{\varepsilon}) \le \phi(y) - \phi(x_{\varepsilon}) + \varepsilon^3,$$

for every $y \in \overline{B}_{R_0}(x_0)$.

Evaluating the DPP at $x = x_{\varepsilon}$ we obtain

$$0 \leq \sup_{A} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} \left(\phi(x_{\varepsilon} + v\varepsilon) - \phi(x) \right) d\sigma(v) \right\} + K\varepsilon^{2}.$$

Since $\phi \in C^2$ performing a second order Taylor expansion and dividing by ε^2 we obtain

$$0 \leq \sup_{A} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} \left(\frac{1}{\varepsilon} \langle \nabla \phi(x_{\varepsilon}), v \rangle + \frac{1}{2} \langle D^{2} \phi(x_{\varepsilon})v, v \rangle + o(1) \right) d\sigma(v) \right\} + K.$$
(23)

Now, since we assumed that $\nabla \phi(x_0) \neq 0$ and $x_{\varepsilon} \to x_0$ we also have $\nabla \phi(x_{\varepsilon}) \neq 0$. Then, after a rotation we can assume that $\nabla \phi(x_{\varepsilon}) = ce_N$ for ε small enough and $\nabla \phi(x_0) = ce_N$. Next we choose $B^* = \{v : \langle e_N, v \rangle \leq \theta_{\varepsilon}\}$ and we arrive to

$$0 \leq \sup_{A} \left\{ \frac{1}{\sigma(A \cap B^*)} \int_{A \cap B^*} \left(\frac{1}{\varepsilon} \langle \nabla \phi(x_{\varepsilon}), v \rangle + \frac{1}{2} \langle D^2 \phi(x_{\varepsilon})v, v \rangle + o(1) \right) d\sigma(v) \right\} + K.$$
(24)

From this inequality we get that for every A with $\sigma(A) \geq \frac{1}{2}\sigma(S^{N-1}) + \eta_{\varepsilon}$, we have

$$0 \ge \frac{1}{\sigma(A \cap B^*)} \int_{A \cap B^*} \frac{1}{\varepsilon} \langle \nabla \phi(x_{\varepsilon}), v \rangle d\sigma(v) \ge -C$$

for some constant C independent of ε .

Using the first inequality, we get

$$0 \leq \sup_{A} \left\{ \frac{1}{\sigma(A \cap B^*)} \int_{A \cap B^*} \left(\frac{1}{2} \langle D^2 \phi(x_{\varepsilon}) v, v \rangle + o(1) \right) d\sigma(v) \right\} + K.$$
(25)

From the geometric measure lemma (Lemma 2.5) we obtain that

$$\frac{1}{\sigma(A\cap B^*)} \int_{A\cap B^*} \frac{1}{2} \langle D^2 \phi(x_{\varepsilon})v, v \rangle d\sigma(v) \to \frac{1}{\mu(\langle e_N, v \rangle = 0)} \int_{\{\langle e_N, v \rangle = 0\}} \frac{1}{2} \langle D^2 \phi(x_0)v, v \rangle d\mu(v).$$

Hence, passing yo the limit as $\varepsilon \to 0$ we arrive to

$$0 \leq \left\{ \frac{1}{\mu(\langle e_N, v \rangle = 0)} \int_{\{\langle e_N, v \rangle = 0\}} \frac{1}{2} \langle D^2 \phi(x_0) v, v \rangle d\mu(v) \right\} + K.$$

$$(26)$$

Now, we just observe that,

$$\frac{1}{\mu(\langle e_N, v \rangle = 0)} \int_{\langle e_N, v \rangle = 0} \frac{1}{2} \langle D^2 \phi(x) v, v \rangle d\mu(v)$$

= $\sum_{i,j=1}^{N-1} \frac{1}{\mu(\langle e_N, v \rangle = 0)} \int_{\langle e_N, v \rangle = 0} \frac{1}{2} \frac{\partial^2 \phi}{\partial v_i \partial v_j}(x) v_i, v_j d\mu(v)$
= $C \sum_{i=1}^{N-1} \frac{\partial^2 \phi}{\partial v_i^2}(x)$

with

$$C = \frac{1}{2} \oint_{\langle e_N, v \rangle = 0} (v_1)^2 d\mu(v).$$

Then, using that we assumed that $\nabla \phi(x_0)$ points in the direction of e_N and that we chose K = C we arrive to

$$0 \le \Delta \phi(x_0) \mid_{\langle \nabla \phi(x_0), v \rangle = 0} + 1, \tag{27}$$

as we wanted to show.

Thus, since $\overline{u}(x) \leq 0$ for $\partial \Omega_0$ (see Proposition 4.2), we obtain that \overline{u} is viscosity subsolution to (2).

Now, we assume that $\nabla \phi(x_0) = 0$. We want to prove that

$$\Delta \phi(x_0) - \langle D^2 \phi(x_0)\eta, \eta \rangle \ge -1,$$

for some vector η with $|\eta| \leq 1$.

As before, we have that for every there exists a sequence $x_{\varepsilon} \in \overline{B}_{R_0}(x_0)$ such that

$$\phi(x_{\varepsilon}) - u^{\varepsilon}(x_{\varepsilon}) - \varepsilon^{3} \leq \inf_{\overline{B}_{R_{0}}(x_{0})} \{\phi(x) - u^{\varepsilon}(x)\} \leq \phi(x) - u^{\varepsilon}(x),$$

and x_{ε} converges to x_0 . Then, as before, we arrive to

$$0 \leq \sup_{A} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} \left(\phi \left(x_{\varepsilon} + v \varepsilon \right) - \phi(x) \right) d\sigma(v) \right\} + k \varepsilon^{2}.$$

Since $\phi \in C^2$ performing a second order Taylor expansion and dividing by ε^2 we obtain

$$0 \leq \sup_{A} \inf_{B} \left\{ \frac{1}{\sigma(A \cap B)} \int_{A \cap B} \frac{1}{\varepsilon} \langle \nabla \phi(x_{\varepsilon}), v \rangle + \frac{1}{2} \langle D^{2} \phi(x_{\varepsilon})v, v \rangle + o(1) d\sigma(v) \right\} + K.$$
(28)

Now, we can argue as follows, given A we choose B as the symmetrized set of A, B = -A. Therefore, we have that $A \cap B = A \cap (-A)$ is symmetric and hence

$$\frac{1}{\sigma(A \cap B)} \int_{A \cap B} \frac{1}{\varepsilon} \langle \nabla \phi(x_{\varepsilon}), v \rangle = 0$$

and

$$\frac{1}{\sigma(A\cap B)} \int_{A\cap B} \frac{1}{2} \langle D^2 \phi(x_{\varepsilon})v, v \rangle) d\sigma(v) = \sum_{i=1}^{N-1} \frac{1}{\sigma(A\cap B)} \int_{A\cap B} \frac{1}{2} \frac{\partial^2 \phi}{\partial v_i^2}(x_{\varepsilon}) d\sigma(v)$$

Hence, we arrive to

$$0 \leq \sup_{A} \sum_{i=1}^{N-1} \frac{1}{\sigma(A \cap B)} \int_{A \cap B} \frac{1}{2} \frac{\partial^2 \phi}{\partial v_i^2}(x_{\varepsilon}) d\sigma(v) + K, \tag{29}$$

and then we obtain

$$\Delta \phi(x_0) - \langle D^2 \phi(x_0) \eta, \eta \rangle \geq -1.$$

To show that \underline{u} is a supersolution, the structure of the proof remains basically the same (with the appropriate changes in the inequalities).

Thus, since $\underline{u}(x) \geq 0$ for $\partial \Omega_0$ (see Proposition 4.2), we obtain that \underline{u} is a viscosity supersolution to (2).

Our aim is to establish the convergence of the value functions of the game, u^{ε} , to the solution to the elliptic problem (2). Having proved that the half relaxed limits satisfy that \overline{u} is a viscosity subsolution and \underline{u} is a viscosity supersolution to (2), our next goal is to prove that \overline{u} and \underline{u} coincide. We will achieve this using the Comparison Principle, Theorem 2.4.

Proof of Theorem 1.3. Consider the upper semicontinuous function \overline{u} and lower semicontinuous function \underline{u} defined by (19).

On the one hand, by definition, we have that $\underline{u}(x) \leq \overline{u}(x)$ for all $x \in \Omega_0$.

By Proposition 4.3, \overline{u} is subsolution and \underline{u} is a supersolution of problem (2). Then, $\underline{u} \geq \overline{u}$ in Ω_0 due to the Comparison Principle, Theorem 2.4. Therefore, we conclude that

$$\overline{u} \equiv \underline{u}$$

which implies that $u := \overline{u} = \underline{u}$ is a continuous function in $\overline{\Omega}_0$. Moreover, u is the unique solution to the elliptic problem (2) and u^{ε} converges to u pointwise in $\overline{\Omega}_0$. In addition, from these convergences we get that the convergence is uniform $\overline{\Omega}_0$, see [19].

Irene Gonzálvez, Alfredo Miranda, Julio D. Rossi and Jorge Ruiz-Cases

4.4 The positivity sets

Finally, recall that we want to study the positivity sets of u^{ε} and u,

$$\Omega_t^{\varepsilon} := \{x : u^{\varepsilon}(x) > t\} \text{ and } \Omega_t := \{x : u(x) > t\}.$$
(30)

Next, as a consequence of the convergence of u^{ε} to u, we will prove that for each fixed t > 0,

$$\Omega_t \subset \liminf_{\varepsilon \to 0} \Omega_t^\varepsilon \subset \limsup_{\varepsilon \to 0} \Omega_t^\varepsilon \subset \overline{\Omega}_t.$$

Proof of Corollary 1.4. Fix t > 0. We start proving that

$$\Omega_t \subset \liminf_{\varepsilon \to 0} \Omega_t^\varepsilon.$$

Suppose that $x \in \Omega_t$. Since

$$\liminf_{\varepsilon \to 0} \Omega_t^{\varepsilon} = \bigcup_{\varepsilon_0 > 0} \bigcap_{0 < \varepsilon \le \varepsilon_0} \Omega_t^{\varepsilon},$$

we want to prove that there exists $\varepsilon_0 > 0$ such that $x \in \Omega_t^{\varepsilon}$ for all $0 < \varepsilon \leq \varepsilon_0$.

Using that $x \in \Omega_t$, we get that there is $\mu > 0$ such that $u(x) \ge t + \mu$. By Theorem 1.3, $u^{\varepsilon} \to u$. Then, there is $\varepsilon_0 > 0$ such that $u^{\varepsilon}(x) \ge t + \mu/2$ for each $0 < \varepsilon \le \varepsilon_0$. Thus, $x \in \Omega_t^{\varepsilon}$ for all $0 < \varepsilon \le \varepsilon_0$.

On the other hand, suppose that $x \notin \overline{\Omega_t}$. Our aim is to prove that

$$x \not\in \limsup_{\varepsilon \to 0} \Omega_t^\varepsilon.$$

$$\limsup_{\varepsilon \to 0} \Omega_t^{\varepsilon} = \bigcap_{\varepsilon_0 > 0} \bigcup_{0 < \varepsilon \le \varepsilon_0} \Omega_t^{\varepsilon},$$

the above is equivalent to prove that there exists $\varepsilon_0 > 0$ for which $x_0 \notin \Omega_t^{\varepsilon}$ for all $0 < \varepsilon \leq \varepsilon_0$. Due to the fact that $x \notin \overline{\Omega_t}$, we get that there exists $\mu > 0$ such that $u(x,t) \leq -\mu$. Using again the convergence of u^{ε} to u, we obtain that there exists $\varepsilon_0 > 0$ such that $u^{\varepsilon}(x) \leq t - \mu/2$ for every $0 < \varepsilon \leq \varepsilon_0$, that is, for each $0 < \varepsilon \leq \varepsilon_0$, $x \notin \Omega_t^{\varepsilon}$.

Acknowledgments

I. Gonzálvez and J. Ruiz-Cases were supported by grants RED2022-134784-T and CEX2023-001347-S. I. Gonzálvez was also supported by grant PID2019-110712GB-I00. J. Ruiz-Cases was also supported by PID2023-146931NB-I00, all of them funded by MI-CIU/AEI/10.13039/501100011033 (Spain). A. Miranda and J. D. Rossi were partially supported by CONICET PIP GI No 11220150100036CO (Argentina) and by UBACyT 20020160100155BA (Argentina).

References

- [1] L. Almeida, A. Chambolle, and M. Novaga. Mean curvature flow with obstacles. Ann. Inst. H. Poincare Anal. Non Lineaire, 29(5):667–681, 2012.
- [2] L. Ambrosio and H. M. Soner. Level set approach to mean curvature flow in arbitrary codimension. J. Diff. Geom., 43:693-737, 1996.
- [3] G. Barles and P.E. Souganidis. Convergence of approximation schemes for fully nonlinear second order equations.. *Asymptot. Anal.*, 4:271–283, 1991.
- [4] P. Blanc and J. D. Rossi. Games for eigenvalues of the Hessian and concave/convex envelopes. J. Math. Pures Appl., 12:192–215, 2019.
- [5] P. Blanc and J. D. Rossi. Game Theory and Partial Differential Equations. *De Gruyter* Series in Nonlinear Analysis and Applications, Vol. 31. 2019.
- [6] K. A. Brakke. *The motion of a surface by its mean curvature*. Princeton University Press, Princeton, N.J., 1978.
- [7] R. Buckdahn, P. Cardaliaguet, and M. Quincampoix. A representation formula for the mean curvature motion. *SIAM J. Math. Anal.*, 33:827–846, 2001.
- [8] M.G. Crandall, H. Ishii, and P.L. Lions. User's guide to viscosity solutions of second order partial differential equations. Bull. Amer. Math. Soc., 27:1–67, 1992.
- [9] J.L. Doob. What is a martingale?. Amer. Math. Monthly, 78(5):451–463, 1971.
- [10] J.L. Doob. Classical Potential Theory and Its Probabilistic Counterpart. Classics in Mathematics, Springer, 2001.
- [11] L.C. Evans and J. Spruck. Motion of level sets by mean curvature. J. Differential Geom., 33(3):635–681, 1991.
- [12] Y.G. Chen, Y. Giga, and S. Goto. Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations. J. Differential Geom., 33(3):749–786, 1991.
- [13] M. Gage and R. S. Hamilton. The heat equation shrinking convex plane curves. J. Differential Geom., 26:69–96, 1986.
- [14] Y. Giga. *Surface evolution equations*. A level set approach. Birkhauser Verlag, Basel, 2006.
- [15] Y. Giga, S. Goto, H. Ishii, and M.H. Sato. Comparison principle and convexity preserving properties for singular degenerate parabolic equations on unbounded domains. *Indiana Univ. Math. J.*, 40(2):443–470, 1991.

Irene Gonzálvez, Alfredo Miranda, Julio D. Rossi and Jorge Ruiz-Cases

- [16] Y. Giga and Q. Liu. A billiard-based game interpretation of the Neumann problem for the curve shortening equation. Adv. Differential Equations, 14(3-4):201-240, 2009.
- [17] Y. Giga, H. Mitake, and H. V. Tran. On asymptotic speed of solutions to level-set mean curvature flow equations with driving and source terms. SIAM J. Math. Anal., 48:3515–3546, 2016.
- [18] Y. Giga, H. V. Tran, and L. Zhang. On obstacle problem for mean curvature flow with driving force. *Geometric flows*, 4:9–29, 2019.
- [19] I. Gonzálvez, A. Miranda, J. D. Rossi, and J. Ruiz-Cases. Finding the convex hull of a set using the flow by minimal curvature with an obstacle. A game theoretical approach. *To appear in Comm. Anal. Geom.*
- [20] M. A. Grayson. The heat equation shrinks embedded plane curves to round points. J. Differential Geom., 26:285–314, 1987.
- [21] R.V. Kohn and S. Serfaty. A deterministic-control-based approach to motion by curvature. Comm. Pure Appl. Math., 59(3):344–407, 2006.
- [22] R.Kohn and S. Serfaty. A deterministic-control-based approach to fully nonlinear parabolic and elliptic equations. *Comm. Pure Appl. Math.*, 10:1298–1350, 2010.
- [23] M. Lewicka. A Course on Tug-of-War Games with Random Noise. Introduction and Basic Constructions. Universitext book series, Springer, 2020.
- [24] M. Lewicka and J. J. Manfredi. The obstacle problem for the *p*-laplacian via optimal stopping of tug-of-war games. *Prob. Theory Rel. Fields*, 167(1-2):349–378, 2017.
- [25] Q. Liu. Fattening and comparison principle for level set equations of mean curvature type. SIAM J. Control Optim., 49:2518–2541, 2011.
- [26] Q. Liu and N. Yamada. An obstacle problem arising in large exponent limit of power mean curvature flow equation. Trans. Amer. Math. Soc., 372:2103–2141, 2019.
- [27] H. Luiro, M. Parviainen, and E. Saksman. Harnack's inequality for p-harmonic functions via stochastic games. Comm. Partial Differential Equations, 38(11):1985–2003, 2013.
- [28] A. P. Maitra and W. D. Sudderth. Discrete gambling and stochastic games. volume 32 of Applications of Mathematics (New York), Springer-Verlag, New York, 1996.
- [29] J. J. Manfredi, M. Parviainen, and J. D. Rossi. An asymptotic mean value characterization for p-harmonic functions. Proc. Amer. Math. Soc., 138(3):881–889, 2010.
- [30] J. J. Manfredi, M. Parviainen, and J. D. Rossi. On the definition and properties of p-harmonious functions. Ann. Scuola Nor. Sup. Pisa, 11:215–241, 2012.

- [31] J. J. Manfredi, J. D. Rossi, and S. J. Somersille. An obstacle problem for tug-of-war games. Commun. Pure Appl. Math., 14:217–228, 2015.
- [32] G. Mercier. Mean curvature flow with obstacles: a viscosity approach, 2014.
- [33] G. Mercier and M. Novaga. Mean curvature flow with obstacles: Existence, uniqueness and regularity of solutions. *Inter. Free Bound*, 17:399–426, 2015.
- [34] A. Miranda and J. D. Rossi. Games for the two membranes problem. Orbita Math., 1(1):59–10, 2024.
- [35] K. Misu. A game-theoretic approach to the asymptotic behavior of solutions to an obstacle problem for the mean curvature flow equation, 2023.
- [36] Y. Peres, O. Schramm, S. Sheffield, and D. Wilson. Tug-of-war and the infinity Laplacian. J. Amer. Math. Soc., 22:167–210, 2009.
- [37] Y. Peres and S. Sheffield. Tug-of-war with noise: a game theoretic view of the p-Laplacian. Duke Math. J., 145(1):91–120, 2008.
- [38] J. D. Rossi. Tug-of-war games and PDEs. Proc. Royal Soc. Edim., 141A:319–369, 2011.
- [39] D. Williams. *Probability with martingales*. Cambridge University Press, Cambridge, 1991.

Received: May 28, 2025 Accepted for publication: June 23, 2025 Communicated by: Jaqueline Godoy Mesquita, Mariel Sáez Trumper, Rafael Potrie and Tiago Macedo